


## PhysioTherapy Exercises - Meniscus Injury

You may do the first 7 exercises right away. You may do the rest of the exercises when the pain in your knee has decreased.

### Passive knee extension

Do this exercise if you are unable to extend your knee fully. While lying on your back, place a rolled-up towel under the heel of your injured leg so the heel is about 6 inches off the ground. Relax your leg muscles and let gravity slowly straighten your knee. Try to hold this position for 2 minutes. Repeat 3 times. You may feel some discomfort while doing this exercise. Do the exercise



### Standing calf stretch

Stand facing a wall with your hands on the wall at about eye level. Keep your injured leg back with your heel on the floor. Keep the other leg forward with the knee bent. Turn your back foot slightly inward (as if you were pigeon-toed). Slowly lean into the wall until you feel a stretch in the back of your calf. Hold the stretch for 15 to 30 seconds. Return to the starting position. Repeat 3 times. Do this exercise several times each day.

### Heel slide

Sit on a firm surface with your legs straight in front of you. Slowly slide the heel of the foot on your injured side toward your buttock by pulling your knee toward your chest as you slide the heel. Return to the starting position. Do 2 sets of 15.





### Hamstring stretch on wall

Lie on your back with your buttocks close to a doorway. Stretch your uninjured leg straight out in front of you on the floor through the doorway. Raise your injured leg and rest it against the wall next to the door frame. Keep your leg as straight as possible. You should feel a stretch in the back of your thigh. Hold this position for 15 to 30 seconds. Repeat 3 times.

## PhysioTherapy Exercises - Meniscus Injury

### Straight leg raise

Lie on your back with your legs straight out in front of you. Bend the knee on your uninjured side and place the foot flat on the floor. Tighten the thigh muscle on your injured side and lift your leg about 8 inches off the floor. Keep your leg straight and your thigh muscle tight. Slowly lower your leg back down to the floor. Do 2 sets of 15.



### Prone hip extension

Lie on your stomach with your legs straight out behind you. Fold your arms under your head and rest your head on your arms. Draw your belly button in towards your spine and tighten your abdominal muscles. Tighten the buttocks and thigh muscles of the leg on your injured side and lift the leg off the floor about 8 inches. Keep your leg straight. Hold for 5 seconds. Then lower your leg and relax. Do 2 sets of 15. sets of 15.

### Clam exercise

Lie on your uninjured side with your hips and knees bent and feet together. Slowly raise your top leg toward the ceiling while keeping your heels touching each other. Hold for 2 seconds and lower slowly. Do 2 sets of 15 repetitions.



### Wall squat with a ball


Stand with your back, shoulders, and head against a wall. Look straight ahead. Keep your shoulders relaxed and your feet 3 feet (90 centimeters) from the wall and shoulder's width apart. Place a soccer or basketball-sized ball behind your back. Keeping your back against the wall, slowly squat down to a 45-degree angle. Your thighs will not yet be parallel to the floor. Hold this position for 10 seconds and then slowly slide back up the wall. Repeat 10 times. Build up to 2 sets of 15.

## PhysioTherapy Exercises - Wobble Board Exercises

Stand on a wobble board with your feet shoulder-width apart. When you have mastered the wobble exercises standing on both legs, try repeating them while standing on just your injured leg. After you are able to do these exercises on one leg, try to do them with your eyes closed. Make sure you have something nearby to support you in case you lose your balance.


### Exercise A

Rock the board forwards and backwards 30 times, then side to side 30 times. Hold on to a chair if you need support.



### Exercise C

Balance on the wobble board for as long as you can without letting the edges touch the floor. Try to do this for 2 minutes without touching the floor.



### Exercise D

Rotate the wobble board in clockwise and counterclockwise circles, but do not let the edge of the board touch the floor.

## PhysioTherapy Exercises - Knee Stabilization

Wrap a piece of elastic tubing around the ankle of your uninjured leg. Tie a knot in the other end of the tubing and close it in a door at about ankle height. Hold onto a chair if you need help balancing. This exercise can be made more challenging by standing on a firm pillow or foam mat while you move the leg with tubing.

### Exercise A

Stand facing the door on the leg without tubing (your injured leg) and bend your knee slightly, keeping your




### Exercise B

Turn 90 degrees so the leg without tubing is closest to the door. Move the leg with tubing away from your body. Do 2

### Exercise C

Turn 90 degrees again so your back is to the door. Move the leg with tubing straight out in front of you.



### Exercise D

Turn your body 90 degrees again so the leg with tubing is closest to the door. Move the leg with tubing across your body. Do 2 sets of 15